

学校编码：10384

学号：21620101152381

厦 门 大 学

硕 士 学 位 论 文

三株TNF- α 拮抗活性菌株的次级代谢产物研究

Studies on the Secondary Metabolites of
three Strains with TNF- α Inhibitory
Activity

楼盈凯

指导教师：徐庆妍

专业名称：微生物学

答辩日期：2013年6月

厦门大学学位论文原创性声明

本人呈交的学位论文是本人在导师指导下，独立完成的研究成果。本人在论文写作中参考其他个人或集体已经发表的研究成果，均在文中以适当方式明确标明，并符合法律规范和《厦门大学研究生学术活动规范(试行)》。

另外，该学位论文为()课题(组)的研究成果，获得()课题(组)经费或实验室的资助，在()实验室完成。(请在以上括号内填写课题或课题组负责人或实验室名称，未有此项声明内容的，可以不作特别声明。)

声明人(签名)：

年 月 日

厦门大学学位论文著作权使用声明

本人同意厦门大学根据《中华人民共和国学位条例暂行实施办法》等规定保留和使用此学位论文，并向主管部门或其指定机构送交学位论文(包括纸质版和电子版)，允许学位论文进入厦门大学图书馆及其数据库被查阅、借阅。本人同意厦门大学将学位论文加入全国博士、硕士学位论文共建单位数据库进行检索，将学位论文的标题和摘要汇编出版，采用影印、缩印或者其它方式合理复制学位论文。

本学位论文属于：

()1. 经厦门大学保密委员会审查核定的保密学位论文，于
年 月 日解密，解密后适用上述授权。

()2. 不保密，适用上述授权。

(请在以上相应括号内打“√”或填上相应内容。保密学位论文应是已经厦门大学保密委员会审定过的学位论文，未经厦门大学保密委员会审定的学位论文均为公开学位论文。此声明栏不填写的，默认为公开学位论文，均适用上述授权。)

声明人(签名)：

年 月 日

摘 要

肿瘤坏死因子- α (TNF- α) 是巨噬细胞或单核细胞活化产生的一种细胞因子, 在炎症反应、肿瘤免疫、细胞死亡、细胞增殖分化等多种生理和病理过程中发挥重要作用。适量的TNF- α 介导的免疫反应对机体具有保护作用, 但过量表达的TNF- α 与其炎症因子一起作用是导致多种病理损伤, 如自身免疫性疾病, 感染性疾病等的原因[], 所以寻找小分子TNF- α 抑制剂, 阻断其过量产生或下游信号通路的传递, 具有非常重要的临床意义。

寻找具有生物活性的新天然产物是发现新药先导化合物的主要途径, 许多临床常用的药物均来自微生物次级代谢产物。而来自深海、极地、盐湖等极端环境中的菌株资源及植物内生真菌资源能够产生种类繁多、结构新颖、活性多样的次级代谢产物, 是获得新结构和新活性天然产物的重要来源。

本论文对两株分离自特殊生境的有TNF- α 拮抗活性的真菌及一株放线菌的次级代谢产物进行了研究, 共分离鉴定了32个化合物, 其中5个为新化合物。

从海藻内生真菌*Aspergillus* sp. AF044固体培养基发酵提取物中分离鉴定了13个化合物, 包括7个生物碱类化合物 (af7, af-8-8', af1023, af1013, af1024, af-8-8, af1210a), 4个苯甲酸类衍生物 (af1, af8, af-91, af1023a), 2个内酯类化合物 (af23e, af422c3), 其中af7、af1024和af1210a为新化合物。

从海洋小单孢菌*Micromonospora* sp. FXY256固体培养基发酵提取物中分离鉴定了4个化合物, 包括2个呋喃酮类化合物 (XY-1-3-1, F-4-4'), 1个内酰胺类 (XY-3), 1个苯丙烯酸 (XY-2)。

从喜树内生真菌*Botryosphaeria* sp. NXG-06固体培养基发酵提取物中分离鉴定了15个化合物, 包括1个呋喃酮类化合物 (M-1-6-10), 5个倍半萜类化合物 (M4-1-2, M5-4-1a, M5-4-1c, M5-4-1f, af11), 2个酮类化合物 (M5-4-1d, N-2-7), 2个二倍半萜类化合物 (M6'-1, N8), 4个环二肽类化合物 (N-2-1, N-2-6b, A-3a-1, A-1-2c-3), 1个环烃类化合物 (m12-1), 其中af11和m12-1为新化合物。

对上述化合物在L929细胞模型上进行拮抗TNF- α 活性筛选，结果显示化合物N-2-1具有一定拮抗活性。

本论文的研究结果表明：海洋微生物蕴藏着丰富的次级代谢产物资源，能够产生结构新颖、具有潜在生物活性的化合物；从天然产物中寻找TNF- α 的小分子抑制剂是可行的。

关键词：次级代谢产物；TNF- α ；真菌；放线菌

厦门大学博硕士论文摘要库

Abstract

Tumor necrosis factor- α (TNF- α) is a proinflammatory cytokine produced by macrophages and neutrophils, it plays an important role in diverse cellular events such as inflammation, tumorigenesis, cell death and differentiation. Appropriate amount of TNF- α mediates protective immune reaction for the host, but overexpression of TNF- α is involved in systemic inflammation and many other acute phase reactions, so blocking the effect of TNF- α has been proved efficient for treating these diseases.

Bioactive natural products are main resource for discovering new drugs. Many important clinical medicines are microbial secondary metabolites. Microorganisms isolated from special living environment, such as deep-sea, polar region, saline and endophytic fungi, et al, can produce numerous metabolites with novel and varied skeletons or with unique bioactivities. In this study, the secondary metabolites of two fungi and one actinomycete with anti- TNF- α activity were investigated. Thirty- two compounds were isolated and elucidated from these three strains, including five new compounds.

The secondary metabolites of the strain *Spergillus* sp. AF044 with agar medium solid were studied. Thirteen compounds were isolated, including seven alkaloids (af7, af-8-8', af1023, af1013, af1024, af-8-8, af1210a), four formic acids (af1, af8, af-91, af1023a) and two lactones (af23e, af422c3), three new compounds (af7, af1024, af1210a) were obtained.

The secondary metabolites of the strain *Micromonospora* sp. FXY256A with agar medium solid were studied. Four compounds were isolated, including two Furan ketones (XY-1-3-1, F-4-4'), one lactam (XY-3), one cinnamic acid (XY-2).

The secondary metabolites of the strain *Botryosphaeria* sp. NXG-06 with agar medium solid were studied. Fifteen compounds were isolated, including one furan ketone (M1-6-10), five terpenoids (M4-1-2, M5-4-1a, M5-4-1c, M5-4-1f, af11),

two ketones(M5-4-1d,N-2-7), two diketopiperazines(M6'-1,N8), four cyclic dipeptides (N-2-1,N-2-6b,A-3a-1,A-1-2c-3) and one hydrocarbon(m12-1). af11 and m12-1 were new ones.

Compound N-2-1 exhibited moderate TNF- α inhibitory activity in L929 cells pretreatment with TNF- α

Our results indicated that marine actinomycetes an specific habitats fungi can produce novel metabolites. Exploring small-molecule inhibitors from the secondary metabolites of microorganisms is feasible.

Keywords: Secondary metabolites; TNF- α ; Fungi; Actinomycetes

参考资料

- [1] Vasanthi P, Nalini G, Rajasekhar G. Role of tumor necrosis factor-alpha in rheumatoid arthritis: a review[J]. *APLAR Journal of Rheumatology*, 2007, 10(4): 270-274.
- [2] DasGupta S, Murumkar P R, Giridhar R, et al. Current perspective of TACE inhibitors: a review[J]. *Bioorganic & medicinal chemistry*, 2009, 17(2): 444-459.
- [3] Cragg G M, Newman D J, Snader K M. Natural products in drug discovery and development[J]. *Journal of Natural products*, 1997, 60(1): 52-60.
- [4] Newman D J, Cragg G M. Natural Products as Sources of New Drugs over the Last 25 Years [J]. *Journal of natural products*, 2007, 70(3): 461-477.
- [5] Bräse S, Encinas A, Keck J, et al. Chemistry and biology of mycotoxins and related fungal metabolites[J]. *Chemical reviews*, 2009, 109(9): 3903-3990.
- [6] Saleem M, Ali M S, Hussain S, et al. Marine natural products of fungal origin[J]. *Natural product reports*, 2007, 24(5): 1142-1152.
- [7] 王兴红. 内生真菌与天然产物研究进展[J]. *中草药*, 2007, 38(1): 140-143.
- [8] Tracey, MD K J, Cerami, Ph. D A. Tumor necrosis factor: A pleiotropic cytokine and therapeutic target[J]. *Annual review of medicine*, 1994, 45(1): 491-503.
- [9] Goldfeld A E, Strominger J L, Doyle C. Human tumor necrosis factor alpha gene regulation in phorbol ester stimulated T and B cell lines[J]. *The Journal of experimental medicine*, 1991, 174(1): 73-81.
- [10] Lupia E, Montrucchio G, Battaglia E, et al. Role of tumor necrosis factor and platelet activating factor in neoangiogenesis induced by synovial fluids of patients with rheumatoid arthritis[J]. *European journal of immunology*, 1996, 26(8): 1690-1694.
- [11] Aggarwal B B, Kohr W J, Hass P E, et al. Human tumor necrosis factor. Production, purification, and characterization[J]. *Journal of Biological Chemistry*, 1985, 260(4): 2345-2354.
- [12] Chen G, Goeddel D V. TNF-R1 signaling: a beautiful pathway[J]. *Science Signaling*, 2002, 296(5573): 1634-1635.
- [13] Shu H B, Takeuchi M, Goeddel D V. The tumor necrosis factor receptor 2 signal transducers TRAF2 and c-IAP1 are components of the tumor necrosis factor receptor 1 signaling complex[J]. *Proceedings of the National Academy of Sciences*, 1996, 93(24): 13973-13978.
- [14] Reinhard C, Shamon B, Shyamala V, et al. Tumor necrosis factor -induced activation of c-jun N-terminal kinase is mediated by TRAF2[J]. *The EMBO Journal*, 1997, 16(5): 1080-1092.
- [15] May M J, D'Acquisto F, Madge L A, et al. Selective inhibition of NF-kappa B activation by a peptide that blocks the interaction of NEMO with the Ikappa B kinase complex[J]. *Science Signaling*, 2000, 289(5484): 1550-1554.
- [16] Tucker S J, Rae C, Littlejohn A F, et al. Switching leukemia cell phenotype between life and death[J]. *Proceedings of the National Academy of Sciences of the United States of America*, 2004, 101(35): 12940-12945.
- [17] Denecker G, Vercammen D, Steemans M, et al. Death receptor-induced apoptotic and necrotic cell death: differential role of caspases and mitochondria[J]. *Cell death and differentiation*, 2001, 8(8): 829-840.
- [18] DasGupta S, Murumkar P R, Giridhar R, et al. Current perspective of TACE inhibitors: a review[J]. *Bioorganic & medicinal chemistry*, 2009, 17(2): 444-459.
- [19] Lin J, Ziring D, Desai S, et al. TNF blockade in human diseases: an overview of efficacy and safety[J]. *Clinical immunology (Orlando, Fla.)*, 2008, 126(1): 13-30.
- [20] Wong M, Ziring D, Korin Y, et al. TNF blockade in human diseases: mechanisms and future directions[J]. *Clinical immunology (Orlando, Fla.)*, 2008, 126(2): 121-136.
- [21] He M M, Smith A S, Oslob J D, et al. Small-molecule inhibition of TNF -alpha[J]. *Science Signaling*, 2005, 310(5750): 1022-1025.
- [22] Natarajan K, Manna S K, Chaturvedi M M, et al. Protein tyrosine kinase inhibitors block tumor necrosis

factor-induced activation of nuclear factor-kappaB, degradation of IkappaBalpha, nuclear translocation of p65, and subsequent gene expression[J]. Archives of biochemistry and biophysics, 1998, 352(1): 59-70.

[23]Jing Z, Wu Z, Song L I. Recent advances of TNF- inhibitors[J].Chinese Journal of Medicinal Chemistry, 2010, 4: 016.

[24] Ward K W, Proksch J W, Azzarano L M, et al. Preclinical Pharmacokinetics of SB-203580, a potent inhibitor of p38 mitogen-activated protein kinase[J]. Xenobiotica, 2001, 31(11): 783-797.

[25] Berdy J. Bioactive microbial metabolites[J]. The Journal of antibiotics, 2005, 58(1): 1-26.

[26]Newman D J, Cragg G M, Snader K M. Natural products as sources of new drugs over the period 1981-2002[J]. Journal of natural products, 2003, 66(7): 1022-1037.

[27]Newman D J, Cragg G M, Snader K M. The influence of natural products upon drug discovery[J]. Natural product reports, 2000, 17(3): 215-234.

[28]Mishra B B, Tiwari V K. Natural products: an evolving role in future drug discovery[J]. European journal of medicinal chemistry, 2011,46(10): 4769-4807.

[29]Fusetani N. Biotechnological potentials of marine natural products[J]. Journal of Biotechnology, 2008, 136: S519.

[30]Rainer Ebel. Comprehensive Natural Products , Volume 2,2010,Pages 223-262.

[31]Blunt J W, Copp B R, Hu W P, et al. Marine natural products[J]. Natural Product Reports, 2009, 26(2): 170-244.

[32]Blunt J W, Copp B R, Munro M H G, et al. Marine natural products[J].Natural product reports, 2010, 27(2): 165-237.

[33]Blunt J W, Copp B R, Munro M H G, et al. Marine natural products[J].Natural Product Reports, 2011, 28(2): 196-268.

[34]Rateb M E, Ebel R. Secondary metabolites of fungi from marine habitats[J]. Natural Product Reports, 2011, 28(2): 290-344.

[35]Abraham E P, Newton G G. New penicillins, cephalosporin C, and penicillinase[J]. Endeavour, 1961, 20: 92-100.

[36]Imhoff J F, Labes A, Wiese J. Bio-mining the microbial treasures of the ocean: new natural products[J]. Biotechnology advances, 2011,29(5): 468-482.

[37]Xia X, Zhang J, Zhang Y, et al. Pimarane diterpenes from the fungus *Epicoccum* sp. HS-1 associated with *Apostichopus japonicus*[J].Bioorganic & medicinal chemistry letters, 2012, 22(8): 3017-3019.

[38]Gao S S, Li X M, Li C S, et al. Penicisteroids A and B, antifungal and cytotoxic polyoxygenated steroids from the marine alga-derived endophytic fungus *Penicillium chrysogenum* QEN-24S[J]. Bioorganic & medicinal chemistry letters, 2011, 21(10): 2894-2897.

[39]Kasettrathat C, Ngamrojanavanich N, Wiyakrutta S, et al. Cytotoxic and antiplasmodial substances from marine-derived fungi, *Nodulisporium* sp. and CRI247-01[J]. Phytochemistry, 2008, 69(14): 2621-2626.

[40]Gu W, Cueto M, Jensen P R, et al. Microsporins A and B: new histone deacetylase inhibitors from the marine-derived fungus *Microsporum* cf. *gypseum* and the solid-phase synthesis of microsporin A[J]. Tetrahedron, 2007, 63(28):6535-6541.

[41]Du L, Li D, Zhu T, et al. New alkaloids and diterpenes from a deep ocean sediment derived fungus *Penicillium* sp[J]. Tetrahedron, 2009, 65(5): 1033-1039.

[42]Wang F, Fang Y, Zhu T, et al. Seven new prenylated indole diketopiperazine alkaloids from holothurian-derived fungus *Aspergillus fumigatus*[J]. Tetrahedron, 2008, 64(34):7986-7991.

[43]Manivasagan P, Venkatesan J, Sivakumar K, et al. Marine actinobacterial metabolites: Current status and future perspectives[J]. Microbiological Research, 2013 168(6):311-332.

[44]Bull A T, Stach J E M. Marine actinobacteria: new opportunities for natural product search and discovery[J]. Trends in microbiology, 2007,15(11): 491-499.

[45]Lam K S. Discovery of novel metabolites from marine actinomycetes[J].Current opinion in microbiology, 2006, 9(3): 245-251.

- [46] Zotchev S B. Marine actinomycetes as an emerging resource for the drug development pipelines[J]. *Journal of Biotechnology*, 2012, 158(4):168-175.
- [47] Subramani R, Aalbersberg W. Marine actinomycetes: An ongoing source of novel bioactive metabolites[J]. *Microbiological Research*, 2012.
- [48] Jensen P R, Williams P G, Oh D C, et al. Species-specific secondary metabolite production in marine actinomycetes of the genus *Salinispora*[J]. *Applied and environmental microbiology*, 2007, 73(4):1146-1152.
- [49] Maloney K N, MacMillan J B, Kauffman C A, et al. Lodopyridone, a structurally unprecedented alkaloid from a marine actinomycete[J]. *Organic letters*, 2009, 11(23): 5422-5424.
- [50] Lu J, Ma Y, Liang J, et al. Aureolic acids from a marine-derived *Streptomyces* sp. WBF16[J]. *Microbiological research*, 2012, 167(10):590-595.
- [51] Gao X, Lu Y, Xing Y, et al. A novel anticancer and antifungal phenazine derivative from a marine actinomycete BM-17[J]. *Microbiological Research*, 2012, 167(10):616-622.
- [52] Sato S, Iwata F, Yamada S, et al. Usabamycins A – C: New anthramycin-type analogues from a marine-derived actinomycete[J]. *Bioorganic & medicinal chemistry letters*, 2011, 21(23): 7099-7101.
- [53] Schulz B, Boyle C, Draeger S, et al. Endophytic fungi: a source of novel biologically active secondary metabolites[J]. *Mycological Research*, 2002, 106(9): 996-1004.
- [54] Lin Z, Zhu T, Fang Y, et al. Polyketides from *Penicillium* sp. JP-1, an endophytic fungus associated with the mangrove plant *Aegiceras corniculatum*[J]. *Phytochemistry*, 2008, 69(5): 1273-1278.
- [55] Zheng C J, Sun P X, Jin G L, et al. Sesquiterpenoids from *Trichoderma atroviride*, an endophytic fungus in *Cephalotaxus fortunei*[J]. *Fitoterapia*, 2011, 82(7): 1035-1038.
- [56] Davis R A, Carroll A R, Andrews K T, et al. Pestalactams A – C: novel caprolactams from the endophytic fungus *Pestalotiopsis* sp[J]. *Organic & biomolecular chemistry*, 2010, 8(8): 1785-1790.
- [57] Chomcheon P, Wiyakrutta S, Sriubolmas N, et al. Aromatase inhibitory, radical scavenging, and antioxidant activities of depsidones and diaryl ethers from the endophytic fungus *Corynespora cassicola* L36[J]. *Phytochemistry*, 2009, 70(3): 407-413.
- [58] Niu X, Xing Wei, Li W, et al. Isofraxidin exhibited anti-inflammatory effects in vivo and inhibited TNF-production in LPS-induced mouse peritoneal macrophages in vitro via the MAPK pathway[J]. *International immunopharmacology*, 2012, 14(2): 164-171.
- [59] Yang L M, Chang S F, Lin W K, et al. Oxygenated compounds from the bioconversion of isostevic acid and their inhibition of TNF- and COX-2 expressions in LPS-stimulated RAW 264.7 cells[J]. *Phytochemistry*, 2012, 75: 90-98.
- [60] Nam S J, Gaudêncio S P, Kauffman C A, et al. Fijiolides A and B, inhibitors of TNF- α -induced NF- κ B activation, from a marine-derived sediment bacterium of the genus *Nocardiopsis*[J]. *Journal of natural products*, 2010, 73(6): 1080-1086.
- [61] Solis P N, Wright C W, Anderson M M, et al. A microwell cytotoxicity assay using *Artemia salina* (brine shrimp)[J]. *Planta medica*, 2007, 59(03): 250-252.
- [62] 鄂征. 细胞培养与分子细胞学技术[M]. 北京出版社, 1995
- [63] Kito K, Ookura R, Yoshida S, et al. Pentaketides relating to aspinonene and dihydroaspyrone from a marine-derived fungus, *Aspergillus ostianus*[J]. *Journal of natural products*, 2007, 70(12): 2022-2025.
- [64] Boutibonnes P. Antibacterial activity of some mycotoxins. [J]. *IRCS Med Sci*, 1980, 8:850-851.
- [65] Tasso G. C. Montenegro, Felipe A. R. Rodrigues, Paula C. Jimenez, Alysson L. Anelin, et al. Cytotoxic Activity of Fungal Strains Isolated from the Ascidian *Eudistoma vancouveri*. *Chemistry & Biodiversity*, 2012, 9:2203-2209.
- [66] Zhang D, Yang X, Kang J S, et al. Circumdatin I, a new ultraviolet-A protecting benzodiazepine alkaloid from a marine isolate of the fungus *Exophiala*[J]. *The Journal of Antibiotics*, 2008, 61(1): 40-42.
- [67] Dai J R, Carté B K, Sidebottom P J, et al. Circumdatin G, a new alkaloid from the fungus *Aspergillus ochraceus*[J]. *Journal of natural products*, 2001, 64(1): 125-126.
- [68] Venkatasubbaiah P, Van Dyke C G, Chilton W S. Phytotoxic metabolites of *Phoma sorghina*, a new foliar

pathogen of pokeweed[J]. *Mycologia*,1992: 715-723.

[69]Ohtake Y, Naito A, Hasegawa H, et al. Novel vasopressin V2 receptor-selective antagonists, pyrrolo [2, 1-a] quinoxaline and pyrrolo [2, 1-c][1, 4] benzodiazepine derivatives[J]. *Bioorganic & medicinal chemistry*, 1999, 7(6): 1247-1254.

[70] Araujo A C, Rauter A P, Nicotra F, et al. Sugar-Based Enantiomeric and Conformationally Constrained Pyrrolo [2, 1-c][1,4]-Benzodiazepines as Potential GABAA Ligands[J]. *Journal of medicinal chemistry*, 2011, 54(5): 1266-1275.

[71]Finefield J M, Kato H, Greshock T J, et al. Biosynthetic Studies of the Notoamides: Isotopic Synthesis of Stephacidin A and Incorporation into Notoamide B and Sclerotiamide[J]. *Organic letters*, 2011, 13(15):3802-3805.

[72]Zhu F, Chen G, Chen X, et al. Aspergicin, a new antibacterial alkaloid produced by mixed fermentation of two marine-derived mangrove epiphytic fungi[J]. *Chemistry of Natural Compounds*, 2011, 47(5):767-769.

[73]Abraham R J, Bernstein H J. The analysis of nuclear magnetic resonance spectra: V. the analysis of deceptively simple spectra [J]. *Canadian Journal of Chemistry*, 1961, 39(1): 216-230.

[74]Duenas M, Fernandez D, Hernandez T, et al. Bioactive phenolic compounds of cowpeas (*Vigna sinensis* L). Modifications by fermentation with natural microflora and with *Lactobacillus plantarum* ATCC 14917[J]. *Journal of the Science of Food and Agriculture*, 2005, 85(2): 297-304.

[75] Duenas M, Fernandez D, Hernandez T, et al. Bioactive phenolic compounds of cowpeas (*Vigna sinensis* L). Modifications by fermentation with natural microflora and with *Lactobacillus plantarum* ATCC 14917[J]. *Journal of the Science of Food and Agriculture*,2005, 85(2):297-304.

[76]Mroczek J E, Rainey W T. Identification and biochemical significance of substitutedfurans in human urine[J]. *Clinical Chemistry*, 1972,18(8): 821-828.

[77]Yuan L, Zhao P J, Ma J, et al. Labdane and Tetranorlabdane Diterpenoids from *Botryosphaeria* sp. MHF, an Endophytic fungus of *Maytenus hookeri*[J]. *Helvetica Chimica Acta*, 2009, 92(6): 1118-1125.

[78] Barrero A F, Arseniyadis S, Quilez del Moral J F, et al. First synthesis of the antifungal oidiolactone C from trans-communic acid: cytotoxic and antimicrobial activity in podolactone-related compounds[J]. *The Journal of Organic Chemistry*, 2002, 67(8): 2501-2508.

[79] Barrero A F, Sánchez J F, Elmerabet J, et al. Enantiospecific syntheses of the potent bioactives nagilactone F and the mould metabolite LL-Z1271 an evaluation of their allelopathic potential[J]. *Tetrahedron*, 1999, 55(23): 7289-7304.

[80] Ichikawa K, Hirai H, Ishiguro M, et al. Cytokine production inhibitors produced by a fungus, *Oidiodendron griseum*[J]. *The Journal of antibiotics*, 2001, 54(9): 697-702.

[81]Ebringer L, Balan J, Catar G, et al. Effect of trypanin on *Toxoplasma gondii* in tissue culture and in mice[J]. *Experimental parasitology*, 1965, 16: 182-189.

[82] Sato M, Kakisawa H. Structures of three new C16 terpenoids from an *Acrostalagmus* fungus[J]. *Journal of the Chemical Society, Perkin Transactions 1*, 1976 (22): 2407-2413.

[83]T. Hosoe, K. Nozawa, T. C. Lumley, R. S. Currah, et al. Tetranordi terpene Lactones, Potent Antifungal Antibiotics for Human Pathogenic Yeasts, from a Unique Species of *Oidiodendron*. *Chem. Pharm. Bull.*, 1999,47(11): 1591-1597.

[84]Logrieco A, Moretti A, Fornelli F, et al. Fusaproliferin production by *Fusarium subglutinans* and its toxicity to *Artemia salina*, SF-9 insect cells, and IARC/LCL 171 human B lymphocytes[J]. *Applied and environmental microbiology*, 1996, 62(9): 3378-3384.

[85]Selvakumar S, Sivasankaran D, Singh V K. Enantioselective Henry reaction catalyzed by C2-symmetric chiral diamine – copper (II) complex[J]. *Organic & Biomolecular Chemistry*,2009, 7(15): 3156-3162.

[86] Rudi A, Kashman Y, Benayahu Y, et al. Amino acid derivatives from the marine sponge *Jaspis digonoxea*[J]. *Journal of natural products*, 1994, 57(6): 829-832.

[87]Yan S, Li S, Wu W, et al. Terpenoid and Phenolic Metabolites from the Fungus *Xylaria* sp. Associated with Termite Nests[J]. *Chemistry & biodiversity*, 2011, 8(9): 1689-1700.

- [88]Fujimoto Y, Yokoyama E, Takahashi T, et al. Studies on the metabolites of *Penicillium diversum* var. *aureum*. I[J]. Chemical & pharmaceutical bulletin, 1986, 34(4): 1497-1500.
- [89]Oka M, Iimura S, Tenmyo O, et al. Terpestacin, a new syncytium formation inhibitor from *Arthrinium* sp[J]. The Journal of antibiotics, 1993, 46(3): 367-373.
- [90]Qi S H, Xu Y, Gao J, et al. Antibacterial and antilarval compounds from marine bacterium *Pseudomonas rhizosphaerae*[J]. Annals of microbiology, 2009, 59(2): 229-233.
- [91]Shigemori H, Tenma M, Shimazaki K, et al. Three New Metabolites from the Marine Yeast *Aureobasidium pullulans*[J]. Journal of natural products, 1998, 61(5): 696-698.
- [92]Wang F W. Bioactive metabolites from *Guignardia* sp., an endophytic fungus residing in *Undaria pinnatifida*[J]. Chinese journal of natural medicines, 2012, 10(1): 72-76.
- [93]Liu H, Chen J, Deng Z W, et al. Secondary metabolites from marine derived fungus *Penicillium terrestre*[J]. (2010)482-486.
- [94]López-Gresa M P, González M C, Primo J, et al. Circumdatin H, a new inhibitor of mitochondrial NADH oxidase, from *Aspergillus ochraceus*[J]. The Journal of antibiotics, 2005, 58(6): 416-419.
- [95]Artman G D, Grubbs A W, Williams R M. Concise, asymmetric, stereocontrolled total synthesis of stephacidins A, B and notoamide B[J]. Journal of the American Chemical Society, 2007, 129(19): 6336-6342.
- [96]Baderschneider B, Winterhalter P. Isolation and characterization of novel benzoates, cinnamates, flavonoids, and lignans from Riesling wine and screening for antioxidant activity[J]. Journal of agricultural and food chemistry, 2001, 49(6): 2788-2798.
- [97]Balan J, Ebringer L, Nemec P, et al. Antiprotozoal antibiotics. ii. isolation and characterization of trypacidin, a new antibiotic, active against *Trypanosoma cruzi* and *Toxoplasma gondii*[J]. The Journal of antibiotics, 1963, 16: 157-60.
- [98]Balan J, Kjaer A, Kovács S, et al. The structure of trypacidin[J]. Acta chemica Scandinavica, 1965, 19: 528-30.
- [99]鲁春华, 黄耀坚, 沈月毛. Secondary Metabolites of *Aspergillus fumigatus* var. *fumigatus*[J]. 中国天然药物, 2005, 3(5): 269-271.
- [100]Jung H J, Lee H B, Kim C J, et al. Anti-angiogenic Activity of Terpestacin, a Bicyclic Sesterterpene from *Embellisia chlamydospora*[J]. J Antibiot, 2003, 56(5):492-496.
- [101]Ritieni A, Monti S M, Randazzo G, et al. Teratogenic effects of fusaproliferin on chicken embryos[J]. Journal of Agricultural and Food Chemistry, 1997, 45(8): 3039-3043.
- [102]Degrazi G, Aguilar C, Bosco M, et al. Plant growth-promoting *Pseudomonas putida* WCS358 produces and secretes four cyclic dipeptides: cross-talk with quorum sensing bacterial sensors[J]. Current microbiology, 2002, 45(4): 250-254.
- [103]Fenical W. Chemical studies of marine bacteria: developing a new resource[J]. Chemical Reviews, 1993, 93(5): 1673-1683.
- [104]Cui C B. Tryprostatins A and B, novel mammalian cell cycle inhibitors produced by *Aspergillus fumigatus*[J]. J. Antibiotics, 1995, 48: 1382-1384.
- [105]Tomassini J E, Davies M E, Hastings J C, et al. A novel antiviral agent which inhibits the endonuclease of influenza viruses[J]. Antimicrobial agents and chemotherapy, 1996, 40(5): 1189-1193.
- [106]McClelland K, Milne P J, Lucieto F R, et al. An investigation into the biological activity of the selected histidine containing diketopiperazines cyclo (His Phe) and cyclo (His Tyr)[J]. Journal of pharmacy and pharmacology, 2004, 56(9): 1143-1153.
- [107]Jamie H, Kilian G, Dyason K, et al. The effect of the isomers of cyclo (Trp Pro) on heart and ion channel activity[J]. Journal of pharmacy and pharmacology, 2002, 54(12): 1659-1665.

Degree papers are in the “[Xiamen University Electronic Theses and Dissertations Database](#)”. Full texts are available in the following ways:

1. If your library is a CALIS member libraries, please log on <http://etd.calis.edu.cn/> and submit requests online, or consult the interlibrary loan department in your library.
2. For users of non-CALIS member libraries, please mail to etd@xmu.edu.cn for delivery details.

厦门大学博硕士论文摘要库